

ClinGen Myeloid Malignancy Variant Curation Expert Panel recommendations for germline *RUNX1* variants

Xi Luo,^{1,*} Simone Feurstein,^{2,*} Shruthi Mohan,³ Christopher C. Porter,⁴ Sarah A. Jackson,⁵ Sioban Keel,⁶ Michael Chicka,⁷ Anna L. Brown,⁸ Chimene Kesserwan,⁹ Anupriya Agarwal,¹⁰ Minjie Luo,¹¹ Zejuan Li,^{12,13} Justyne E. Ross,³ Panagiotis Baliakas,¹⁴ Daniel Pineda-Alvarez,¹⁵ Courtney D. DiNardo,¹⁶ Alison A. Bertuch,¹ Nikita Mehta,¹⁷ Tom Vulliamy,¹⁸ Ying Wang,¹⁹ Kim E. Nichols,⁹ Luca Malcovati,²⁰ Michael F. Walsh,²¹ Lesley H. Rawlings,²² Shannon K. McWeeney,²³ Jean Soulier,²⁴ Anna Raimbault,²⁴ Mark J. Routbort,²⁵ Liying Zhang,²⁶ Gabriella Ryan,²⁷ Nancy A. Speck,²⁸ Sharon E. Plon,¹ David Wu,^{29,†} and Lucy A. Godley^{2,†}

¹Department of Pediatrics/Hematology-Oncology, Baylor College of Medicine, Houston, TX; ²Section of Hematology/Oncology and Center for Clinical Cancer Genetics, The University of Chicago, Chicago, IL; ³Department of Genetics, University of North Carolina School of Medicine, Chapel Hill, NC; ⁴Department of Pediatrics, Emory University School of Medicine, Atlanta, GA; ⁵GeneDx, Gaithersburg, MD; ⁶Division of Hematology, Department of Medicine, University of Washington, Seattle, WA; ⁷PreventionGenetics, Marshfield, WI; ⁸Centre for Cancer Biology, SA Pathology & University of South Australia, Adelaide, Australia; ⁹Department of Oncology, St. Jude Children's Research Hospital, Memphis, TN; ¹⁰Knight Cancer Institute, Oregon Health & Science University, Portland, OR; ¹¹Department of Pathology and Laboratory Medicine, The Children's Hospital of Philadelphia, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; ¹²Department of Pathology and Genomic Medicine, Houston Methodist Research Institute and Houston Methodist Hospital, Houston, TX; ¹³Department of Pathology and Laboratory Medicine, Weill Cornell Medical College, New York, NY; ¹⁴Department of Immunology, Genetics, and Pathology, Science for Life Laboratory, Uppsala University, Uppsala, Sweden; ¹⁵Invitae, San Francisco, CA; ¹⁶Department of Leukemia, University of Texas MD Anderson Cancer Center, Houston, TX; ¹⁷Hematopathology Division, Department of Laboratory Medicine & Pathology, Mayo Clinic, Rochester, MN; ¹⁸Blizard Institute, Barts and The London School of Medicine and Dentistry, Queen Mary University London, London, United Kingdom; ¹⁹Department of Pathology and Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, MD; ²⁰Department of Molecular Medicine, University of Pavia and Istituto di Ricovero e Cura a Carattere Scientifico Policlinico S. Matteo Foundation, Pavia, Italy; ²¹Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York, NY; ²²Department of Genetics and Molecular Pathology, SA Pathology, Adelaide, Australia; ²³Division of Biostatistics, Department of Public Health and Preventive Medicine, Oregon Health & Science University, Portland, OR; ²⁴INSERM/CNRS U944/7212, Université de Paris and Hematology Laboratory Assistance Publique-Hôpitaux de Paris, Hôpital Saint-Louis, Paris, France; ²⁵Department of Hematopathology, University of Texas MD Anderson Cancer Center, Houston, TX; ²⁶Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY; ²⁷Department of Scientific Affairs, American Society of Hematology, Washington, DC; ²⁸Department of Cell and Developmental Biology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA; and ²⁹Department of Laboratory Medicine, University of Washington, Seattle, WA

Key Points

- The ClinGen MM-VCEP has specified *RUNX1*-specific curation rules to address gene function, gene-specific domains, and phenotypic criteria.
- *RUNX1*-specific criteria resulted in a reduction in CONF and VUS variants by 33%, emphasizing the need for expert variant curation.

Standardized variant curation is essential for clinical care recommendations for patients with inherited disorders. Clinical Genome Resource (ClinGen) variant curation expert panels are developing disease-associated gene specifications using the 2015 American College of Medical Genetics and Genomics (ACMG) and Association for Molecular Pathology (AMP) guidelines to reduce curation discrepancies. The ClinGen Myeloid Malignancy Variant Curation Expert Panel (MM-VCEP) was created collaboratively between the American Society of Hematology and ClinGen to perform gene- and disease-specific modifications for inherited myeloid malignancies. The MM-VCEP began optimizing ACMG/AMP rules for *RUNX1* because many germline variants have been described in patients with familial platelet disorder with a predisposition to acute myeloid leukemia, characterized by thrombocytopenia, platelet functional/ultrastructural defects, and a predisposition to hematologic malignancies. The 28 ACMG/AMP codes were tailored for *RUNX1* variants by modifying gene/disease specifications, incorporating strength adjustments of existing rules, or both. Key specifications included calculation of minor allele frequency thresholds, formulating a semi-quantitative approach to counting multiple independent variant occurrences, identifying functional domains and mutational hotspots, establishing functional assay thresholds, and characterizing phenotype-specific guidelines. Preliminary rules were tested by using a pilot set of 52 variants; among these, 50 were previously

Submitted 1 July 2019; accepted 24 August 2019. DOI 10.1182/bloodadvances.2019000644.

*X.L. and S.F. contributed equally to this study and are joint first authors.

†D.W. and L.A.G. are joint senior authors.

The full-text version of this article contains a data supplement.

© 2019 by The American Society of Hematology

classified as benign/likely benign, pathogenic/likely pathogenic, variant of unknown significance (VUS), or conflicting interpretations (CONF) in ClinVar. The application of *RUNX1*-specific criteria resulted in a reduction in CONF and VUS variants by 33%, emphasizing the benefit of gene-specific criteria and sharing internal laboratory data.

Introduction

In 2015, the American College of Medical Genetics and Genomics (ACMG) and the Association for Molecular Pathology (AMP) released a landmark document providing guidance on variant classification that has now been adopted by many international diagnostic laboratories. It was designed to have universal applicability to all Mendelian disorders, using several types of weighted and categorized evidence, and it therefore requires significant expertise as well as gene- and disease-specific knowledge to be correctly applied.¹ Variable application of functional and domain-related evidence and inconsistent interpretation and use of the ACMG/AMP criteria are key contributors to incorrect classifications of variants, and significant discrepancies among laboratories highlight the utility of expert guidance.²⁻⁴ A few studies have proposed approaches to one or more aspects of variant interpretation, such as quantitative criteria for cosegregation, use of population databases, adaptation of minor allele frequency (MAF), classes of evidence, and gene-level implications.²⁻⁸ However, due to the unique characteristics of every gene and its disease correlates, along with the variability in the application of classification criteria and evidence interpretation, there is still a lack of comprehensive guidance for variant interpretation.

This need for expert involvement and gene-specific guidance has been addressed by the National Institutes of Health (NIH)-funded Clinical Genome Resource (ClinGen; <https://clinicalgenome.org>), which serves as a body for managing and centralizing clinically relevant genomic knowledge, providing guidance and tools for defining the clinical validity of gene and variant contributions to disease. Several working groups and expert panels were created within ClinGen, including gene- and disease-specific Variant Curation Expert Panels (VCEPs).^{9,10} Moreover, the ClinGen Sequence Variant Interpretation (SVI) Working Group (<https://clinicalgenome.org/working-groups/sequence-variant-interpretation/>) aims to provide general recommendations for the refinement and evolution of the ACMG/AMP guidelines, which are then specialized further by the gene-specific VCEP.

The publicly available ClinVar database (<https://www.ncbi.nlm.nih.gov/clinvar/>), launched in 2013, serves as a valuable centralized resource for documenting the clinical significance of genetic variants submitted by clinical and research laboratories and databases such as OMIM (Online Mendelian Inheritance in Man) and GeneReviews.¹¹ ClinVar uses the ACMG-recommended 5-level scoring system to indicate the level of evidence supporting the assertion of clinical significance of a variant. Human variant data curated by ClinGen expert panels are submitted to ClinVar with a 3-star status (reviewed by expert panel) including a designation that the ClinGen VCEP process has been recognized by the US Food and Drug Administration (FDA).

The general workflow of a VCEP is to define its leadership/membership and scope of focus as well as conflicts of interest.⁹

Once approved, this group develops disease-specific variant classification rules, based on curation of gene-specific features, published literature, and evidence thresholds that are consistent with gene-disease associations. ACMG and AMP have defined 28 codes that address specific evidence, including population data, segregation data, functional data, computational predictions, and allelic data.³ Each code is weighted according to the strength of the evidence: stand-alone, very strong, strong, moderate, or supporting. Codes are also designated as defining the direction of clinical significance: benign (BEN) or pathogenic (PATH). These evidence codes applied to variants are then combined to arrive at a single designation of clinical significance: PATH, likely pathogenic (LPATH), variant of uncertain significance (VUS), likely BEN (LBEN), or BEN. Once preliminary rules are specified, they are pilot tested on a collection of variants with existing assertions of clinical significance, and based on the results of this preliminary testing, the VCEP may adjust some of its rules to optimize variant classification. Once final rules have been approved, they are published and implemented, with VCEP-curated assertions disseminated via the ClinVar database.

A Myeloid Malignancy VCEP (hereafter referred to as MM-VCEP) was formed in 2018 as a collaboration between the American Society of Hematology and ClinGen. The MM-VCEP began adapting the ACMG/AMP framework for *RUNX1* variant classification. Because it was the first germline predisposition syndrome identified for myeloid malignancies, there were many variants already deposited in the ClinVar repository. Germline PATH variants in *RUNX1*, first described in 1999, cause dominantly inherited familial platelet disorder with a predisposition to acute myeloid leukemia (FPD/AML), characterized by mild to moderate thrombocytopenia, functional and ultrastructural platelet defects, and a predisposition to myelodysplastic syndrome (MDS) and AML and less frequently to T-cell acute lymphoblastic leukemia (T-ALL).¹²⁻¹⁴ In 2016, the revision of the World Health Organization classification of myeloid neoplasms and acute leukemia included myeloid malignancies arising from germline PATH variants in *ANKRD26*, *ETV6*, and *RUNX1* in a new category defined as “myeloid neoplasms with germline predisposition and preexisting platelet disorder.”¹⁵ Reported inherited and de novo *RUNX1* variants include missense, nonsense, and splice site single-nucleotide variants (SNVs), small in- or out-of-frame insertions and deletions (indels), as well as copy number variants (CNVs) such as intragenic or whole-gene deletions.¹⁶⁻¹⁸ The prevalence of PATH *RUNX1* germline variants is unknown but presumed to be rare. The disease shows high penetrance with variable expressivity and genotype/phenotype correlation, and the lifetime risk of hematologic malignancies is ~44%, with an average age of onset of 33 years.¹⁹⁻²¹ More than one-half of germline *RUNX1* variants are reported in single probands/families,¹³ leading to a high allelic heterogeneity that restricts the collection of data from segregation analyses and

functional analyses across several affected families. Individuals with a hematologic malignancy are often candidates for hematopoietic stem cell transplantation. The identification of patients with a PATH germline variant in *RUNX1* and its correct classification of the variant are imperative to the selection of potential related donors, among other clinical implications.²²⁻²⁶

Here, we present the *RUNX1*-specific guidelines generated by the MM-VCEP. The MM-VCEP adapted the ACMG/AMP framework for *RUNX1* variant classification with the aim of improving consistency in variant classification and curating *RUNX1* variants for 3-star submission to ClinVar. We used multiple lines of evidence, showing the rationale and data supporting each criterion's modification, and the results from pilot testing the criteria on variants with BEN/LBEN, PATH/LPATH, VUS, and conflicting (CONF) ClinVar assertions. The application of rules for *RUNX1* variant curation will serve as a model for the curation of variants in other genes that also cause inherited myeloid hematologic malignancies, such as *ANKRD26*, *ETV6*, *DDX41*, and *GATA2*. The ClinGen's Web site contains the MM-VCEP variant classification recommendations and any subsequent modifications to these codes over time (<https://www.clinicalgenome.org/affiliation/50034>).

Methods

ClinGen MM-VCEP

The MM-VCEP is sponsored by the American Society of Hematology through its partnership with ClinGen and is described at <https://clinicalgenome.org/affiliation/50034/>. The MM-VCEP team comprises 34 professionals with expertise in key domains and includes clinical geneticists, genetic counselors, hematologists with professional training in genetics, laboratory and research scientists, and variant curation experts. Additional emphasis was placed on global representation, with 22 participating institutions in 6 countries: Australia, France, Italy, Sweden, the United Kingdom, and the United States. The MM-VCEP meets regularly via biweekly teleconferences and corresponds via e-mail on a regular basis. Approval of MM-VCEP is overseen by ClinGen and consists of 4 steps: (1) defining the group/members and scope of the VCEP; (2) developing gene/disease-specific classification rules; (3) optimization of rules using pilot variants; and (4) MM-VCEP approval by ClinGen, implementation of rules in the ClinGen Variant Curation Interface, and submission of curated variants to the ClinVar database. For step two, members were divided into 3 subgroups that focused on the modification of functional/computational/splicing criteria (Team F), population/phenotypic criteria (Team P), and segregation/allelic/de novo criteria (Team S). All members disclosed potential conflicts of interest as required by ClinGen.

ACMG/AMP specifications for *RUNX1*

MM-VCEP members proposed and discussed changes to the existing ACMG/AMP classifications for *RUNX1* germline variants and arrived at consensus decisions via teleconference calls and e-mail. Criteria modifications included gene- or disease-specific modifications, strength-level adjustments, general recommendations, and certain criteria being deemed "not applicable." Publicly available databases, predictive software, and published data obtained from relevant papers were used for criteria specifications. For BA1/BS1 *RUNX1*-specific population MAF, calculations were made assuming Hardy-Weinberg equilibrium using the recently

published Whiffin/Ware online calculator.⁶ Additional efforts included identification of key functional domains and mutational hotspots within *RUNX1*, definition of informative functional assays, and characterization of phenotypic criteria. Recommendations for using ACMG/AMP criteria from the ClinGen's SVI working group were also incorporated.²⁷⁻²⁹ Preliminary and final ACMG/AMP specifications required complete consensus of the MM-VCEP.

Pilot variants

All pilot variants are annotated by using RefSeq IDs NM_001754.4 and NC_000021.9 (GRCh38/hg38). Variants submitted to ClinVar by a variety of clinical laboratories were prioritized for classification. Preliminary rules were refined by interpreting a set of 52 *RUNX1* variants, which were selected to represent the spectrum of variants in *RUNX1*, covering various types of SNVs such as missense, nonsense, splice site, synonymous, and intronic variants; indels such as in-frame duplications and out-of-frame deletions; and CNVs such as intragenic deletions. Similarly, the pilot variants covered a diverse range of classifications in ClinVar, including discrepant assertions (12 BEN/LBEN, 14 VUS, 20 PATH/LPATH, 4 CONF, and 2 with no ClinVar assertions). The variant classification and rules applied were reviewed on conference calls to resolve discrepancies and reach consensus. Basic information regarding individual phenotypes and segregation with disease was obtained from the literature and ClinVar submitters. Statistical approaches for calculations of PS4 are available in the supplemental Methods. Further optimization of rules was performed, and a discussion with the entire MM-VCEP was triggered whenever members disagreed or raised concerns regarding the applicability of a given rule. Curators used ClinGen's Variant Curation Interface (<https://curation.clinicalgenome.org>) to assess and document the applicable rules for each variant. Once the MM-VCEP was approved, the classified *RUNX1* variants with the adapted evidence code framework applied to the variants were submitted to ClinVar and were designated with a 3-star evidence code and FDA recognition flag. The first 52 *RUNX1* variant curations are now available in ClinVar and can be accessed at <https://www.ncbi.nlm.nih.gov/clinvar/submitters/507107/>.

Results

Summary of rule specifications

The final MM-VCEP ACMG/AMP specifications for *RUNX1* were approved by ClinGen and are outlined in Table 1. Six of the original 28 ACMG/AMP criteria had general recommendations on the application of the rule (PM2, PP3, BS4, BP2, BP4, and BP7), 2 required gene- or disease-based specifications (BA1 and BS1), and 2 rules were adjusted in their level of strength (PS1 and PM5). Both gene- or disease-based and strength-level specifications were made to 9 rules (PVS1, PS2, PS3, PS4, PM1, PM4, PM6, PP1, and BS3). Five rules required exceptions for combinations with other rules (PS2, PS3, PM5, PM6, and PP3), and 9 rules were deemed not applicable (PM3, PP2, PP4, PP5, BS2, BP1, BP3, BP5, and BP6). One change to the ACMG/AMP combination of criteria for classification of clinical significance was made in the case of BS1, which can be used as a stand-alone criterion for LBEN classification in the absence of any supporting PATH evidence. The following section highlights the approaches and rationale behind key specifications such as phenotypic criteria, MAF thresholds, and validity of functional assays. Of note, germline material for patients

with FPD/AML or patients with suspected inherited hematologic malignancies cannot include blood or bone marrow from these patients because this is the affected tissue harboring somatic mutations. We recommend using cultured skin fibroblasts as the gold standard, or alternatively DNA from hair roots or cultured mesenchymal stromal cells.^{19,25}

Phenotypic criteria for FPD/AML

FPD/AML is characterized by mild to moderate thrombocytopenia, platelet functional and/or ultrastructural defects, and a predisposition to hematologic malignancies, most often AML and MDS, and less frequently T-ALL (Table 2). The penetrance is high; however, not all individuals carrying the PATH variant display the FPD/AML phenotype. Thrombocytopenia is the most common clinical presentation, followed by hematologic malignancies in ~44% of these patients.¹⁹⁻²¹ The MM-VCEP defined that in order to fit the FPD/AML phenotype, the patient must exhibit at least one of the following phenotypic criteria: (1) mild to moderate thrombocytopenia with normal platelet size and volume in the absence of other causative factors such as autoimmune (eg, antibodies against platelet surface antigens) or drug-related thrombocytopenia³⁰; (2) platelet ultrastructural and/or functional defects, including platelet alpha³¹ or dense granule secretion defects^{30,32,33} or impaired platelet aggregation, particularly in response to collagen and epinephrine^{34,35}; and (3) diagnosis of a hematologic malignancy, most commonly affecting the myeloid lineage causing AML or MDS, less frequently involving the lymphoid lineage and manifesting as T-ALL.^{26,30,36,37} There are rare case reports of patients with germline *RUNX1* variants and mixed myeloproliferative syndromes/MDS such as chronic myelomonocytic leukemia,^{26,38} as well as case reports of patients with B-cell ALL³⁹ and hairy-cell leukemia.⁴⁰

Population data (BA1, BS1, PM2, PS4, PS4_moderate, PS4_supporting, and BP2)

FPD/AML is a rare disorder. The prevalence of the disease-associated *RUNX1* variants is unknown, with an estimated 5515 families worldwide based on a population incidence generated from a survey of centers with FPD/AML patients (A.L.B., written communication, 10 June 2019), which is likely an underestimate of the true prevalence. Among the 3 phenotypic features seen in individuals with germline *RUNX1* variants (Table 2), thrombocytopenia is the most common. We conservatively estimated the prevalence of thrombocytopenia for use in the BA1/BS1 calculations. Most clinical laboratories establish their reference values for platelet counts by measuring samples from at least 120 healthy individuals and identifying the most outlying 5% of observed values. Most often, these outlying observations are split evenly between the ends of the test result distribution in the reference population (2.5% at each end of the distribution), resulting in a 2-sided reference interval.⁴¹ Using this approach, the prevalence of thrombocytopenia can be defined as 1 in 40. The penetrance in families with *RUNX1* germline variant is high to near-complete, with 85% being the lowest penetrance reported to date^{13,19-21} (S.K., written communication, 19 March 2019). Thus far, no founder variants in *RUNX1* have been reported. De novo variants are rare but have been described.¹⁶⁻¹⁸

The MM-VCEP modified BA1 using these conservative assumptions and corresponding values to account for the unknown prevalence and disease contribution of *RUNX1*. To obtain an *RUNX1*-specific population MAF threshold for BA1, we used the

Whiffin/Ware calculator⁶ (<http://cardiodb.org/allelefrequencyapp/>) with a prevalence of 1 in 40, a conservative unascertained penetrance estimate of 85%, an allelic heterogeneity of 100%, and a maximum genetic heterogeneity of 10%. The MM-VCEP also adopted the SVI recommendation that the variant be present in any general continental population dataset with a minimum number of 2000 alleles and the variant present in ≥ 5 alleles.⁴² A 95% confidence interval was used to develop the thresholds. The threshold developed for application of BA1 as a stand-alone criterion is a MAF ≥ 0.0015 (0.15%). For BS1, a maximum genetic heterogeneity contribution of 1% (1 magnitude lower) was used, which led to a range of 0.00015 (0.015%) to 0.0015 (0.15%) for application of BS1. Because we used conservative values for the calculation, we allow a variant to reach an LBEN classification based on BS1 alone if there is no contradictory evidence supporting pathogenicity (as outlined in a recent SVI revision).²⁹ For this work, the gnomAD population database was mostly used, although other databases with a minimum of 2000 alleles are also sufficient. However, we encourage the use of a large dataset such as gnomAD, ExAC, or ESP.

Because most *RUNX1* variants are unique to probands or families,¹³ it was determined that the variant must be completely absent from all population databases to apply PM2. The MM-VCEP tested pilot PATH/LPATH variants with this rule and validated this determination. The MM-VCEP further recommends that the mean coverage of exome and genome sequencing data for *RUNX1* in the population databases used should be at least 20 \times .

Criterion PS4 is based on the significantly higher prevalence of a variant in case cohorts vs control cohorts, which is considered strong evidence for pathogenicity. Ideally, published case-control studies are used as evidence. Given the rarity of FPD/AML, an existing case-control study for *RUNX1* variants could not be identified. The original ACMG/AMP guideline states that the odds ratio (OR), measuring an association between a genotype and phenotype, can be used for Mendelian diseases. Accordingly, in the absence of a published case-control study, the MM-VCEP created a “quasi-case-control study” with the estimated number of probands worldwide and the overall gnomAD population as the control cohort. To apply this code, the proband has to meet at least one of the *RUNX1*-phenotypic criteria (Table 2), and the variant has to be either absent from gnomAD or only present once. This code has a sliding weight scale to account for the number of unrelated probands who meet the *RUNX1*-phenotypic criteria. PS4 is applied with ≥ 4 probands (OR, 100.6), PS4_moderate with 2 to 3 probands (OR, 50.3-75.5), and PS4_supporting with 1 proband (OR, 25.1) (supplemental Table 1).

BP2, supporting evidence for a BEN code, can be applied in the context of autosomal dominant FPD/AML when the variant is found in trans with a known PATH variant. Because there is no evidence in the literature of probands with a homozygous PATH *RUNX1* variant, and lack of Runx1 is embryonically lethal in mice, the MM-VCEP recommends that BP2 also be applied when a variant is found in a confirmed homozygous state in population databases or internal laboratories.^{43,44}

Segregation data (PP1_strong, PP1_moderate, PP1, and BS4)

Segregation with disease (PP1) is used as evidence for pathogenicity, and with increasing number of meioses, a stronger level of

Table 1. MM-VCEP ACMG/AMP specifications for *RUNX1* variants

ACMG/AMP criteria code	Original ACMG/AMP rule summary	Specification	Stand-alone	Very strong	Strong	Moderate	Supporting	Comments
PVS1	Null variant in a gene where LOF is a known mechanism of disease	Gene-specific, strength	NA	Per modified <i>RUNX1</i> PVS1 decision tree for SNVs, indels, and CNVs and table of splicing effects	Same AA change as a previously established PATH variant regardless of nucleotide change	Same AA change as a previously established, LPATH variant regardless of nucleotide change	NA	<i>RUNX1</i> LOF variants are a common mechanism of disease in FPD/AML. Three major isoforms (A, B, and C) are expressed by use of 2 promoters and alternative splicing. C-terminal variants not predicted to undergo NMD are classified as PVS1_strong , deletions of exons 2 and 3, presumably only affecting <i>RUNX1</i> isoform 1C, meet PVS1_moderate
PS1	Same AA change as a previously established PATH variant regardless of nucleotide change	Strength	NA	NA	Same AA change as a previously established, LPATH variant regardless of nucleotide change	Same AA change as a previously established, LPATH variant regardless of nucleotide change	NA	(1) RNA data or agreement in splicing predictors show no splicing effects (SSF and MES predict either increase in canonical splice site score or decrease in canonical splice score by no more than 10% and no putative splice sites are created). (2) The previously established variant must be asserted PATH/LPATH based on MM-VCEP rules for <i>RUNX1</i> before this rule can be applied
PS2	De novo (maternity and paternity confirmed) in a patient with the disease and no family history	Disease-specific, strength	NA	NA	Two or more proven de novo occurrences (maternity and paternity confirmed) in patients with the <i>RUNX1</i> -phenotype	Two or more proven de novo occurrences (maternity and paternity confirmed) in patients with the <i>RUNX1</i> -phenotype	One proven de novo occurrence (maternity and paternity confirmed) in a patient with the <i>RUNX1</i> -phenotype	(1) No family history is defined as: absence of the variant and any of the <i>RUNX1</i> -phenotypic criteria in first- and/or second-degree relatives. (2) The proband must exhibit at least 1 phenotypic FPD/AML criterion. (3) The maximum allowable strength by combining PS2 and PM6 criteria is to apply 1 moderate or 2 supporting rules
PS3	Well-established in vitro or in vivo functional studies supportive of a damaging effect	Gene-specific, strength	NA	NA	Transactivation assays exhibiting altered transactivation (<20% of wt, and/or reduced to levels similar to well-established PATH variants such as R201Q or R166Q) and data from a secondary assay showing altered function. PS3 cannot be applied if the variant meets PVS1 . If the variant meets criteria for PVS1_strong and PS3 , we recommend either applying PVS1_strong and PS3_moderate or upgrading PVS1_strong to PVS1 without applying PS3	Transactivation assays exhibiting altered transactivation (<20% of wt, and/or reduced to levels similar to well-established PATH variants such as R201Q or R166Q) or ≥2 secondary assays exhibiting altered function	Transactivation assays exhibiting enhanced transactivation (>115% of wt)	(1) Transactivation assays should include wt and known PATH controls, as well as coexpression with CBFβ. Promoter sequences of CSF1R (M-CSF-R), PF4, C-FMS, and GZMB, containing consensus <i>RUNX1</i> -binding sites have been used for transactivation assays. (2) The following secondary assays have been performed: EMSA and yeast hybrid assays (decreased DNA-binding affinity), co-IP, FRET, and affinity assays (diminished heterodimerization ability with CBFβ). IF and WB with cell fractionation (abnormal cellular localization), colony-forming assays (reduced colony-forming potential), and

Evidence codes are noted in bold font.

AA, amino acid; co-IP, coimmunoprecipitation; EMSA, electrophoretic mobility shift assay; FRET, fluorescence resonance energy transfer; IF, immunofluorescence; LOF, loss-of-function; MES, MaxEntScan; NA, not applicable; NMD, nonsense-mediated decay; RHD, Runt homology domain; SSF, Splice Site Finder; WB, western blot; wt, wild type.

Table 1. (continued)

ACMG/AMP criteria code	Original ACMG/AMP rule summary	Specification	Stand-alone	Very strong	Strong	Moderate	Supporting	Comments
PS4	The prevalence of the variant in affected individuals is significantly increased compared with the prevalence in control subjects	Disease-specific, strength	NA	NA	Four or more probands meeting <i>RUNX1</i> -phenotypic criteria	Two to 3 probands meeting <i>RUNX1</i> -phenotypic criteria	One proband meeting <i>RUNX1</i> -phenotypic criteria	xenotransplantation experiments (abnormal function of mutant <i>RUNX1</i> in vivo). (3) PS3 can also be applied for evidence of very low or abnormal mRNA/protein expression of the variant allele as a functional consequence of a null variant or incorrect mRNA/protein products
PM1	Located in a mutational hotspot and/or critical and well-established functional domain without BEN variation	Gene-specific, strength	NA	NA	NA	Variant affecting 1 of the other following 13 hotspot residues: R107, K110, A134, R162, R166, S167, R169, G170, K194, T196, D198, R201, R204	Variant affecting 1 of the other AA residues 105-204 within the RHD	The affected individual has to fit at least 1 of the <i>RUNX1</i> -phenotypic criteria and the variant has to be either absent from gnomAD (overall population) or only present once The RHD (AA 77-204) has been established as a highly conserved DNA-binding domain without any BEN variation in ClinVar. No germline PATH variants have been reported in residues in the region (AA 77-104) to date. The AA range under PM1_supporting may be expanded in the future to other parts of the protein if more evidence emerges
PM2	Absent from control subjects	General recommendation	NA	NA	NA	Per original ACMG/AMP guidelines	NA	Variant must be completely absent from all population databases. The mean coverage of <i>RUNX1</i> in the population database used should be at least 20×
PM3	For recessive disorders, detected in trans with a PATH variant	NA	NA	NA	NA	NA	NA	FDP/AML is inherited in an autosomal dominant manner
PM4	Protein length changes due to in-frame deletions/insertions in a nonrepeat region or stop-loss variants	Gene-specific, strength	NA	NA	NA	In-frame deletion/insertion affecting at least 1 of the 13 hotspot residues (R107, K110, A134, R162, R166, S167, R169, G170, K194, T196, D198, R201, R204)	Other in-frame deletion/insertion affecting residues 105-204 within the RHD	See PM1
PM5	Missense change at AA residue where a different missense change determined to be PATH has been seen before	Strength	NA	NA	Missense change at the same residue where ≥ 2 different missense changes have previously been determined to be PATH. PM5_strong cannot be applied together with PM1	Missense change at the same residue where a different missense change has previously been determined to be PATH	Missense change at the same residue where a different missense change has previously been determined to be LPATH	See PS1
PM6	Assumed de novo (but without confirmation of maternity and paternity) in a patient with the disease and no family history	Disease-specific, strength	NA	NA	NA	Four or more assumed de novo occurrences (without confirmation of maternity and paternity) in patients with the <i>RUNX1</i> -phenotype	Two or 3 assumed de novo occurrences (without confirmation of maternity and paternity) in patients with the <i>RUNX1</i> -phenotype	See PS2

Evidence codes are noted in bold font.

AA, amino acid; co-IP, coimmunoprecipitation; EMSA, electrophoretic mobility shift assay; FRET, fluorescence resonance energy transfer; IF, immunofluorescence; LOF, loss-of-function; MES, MaxEntScan; NA, not applicable; NMD, nonsense-mediated decay; RHD, Runt homology domain; SSF, Splice Site Finder; WB, western blot; wt, wild type.

Table 1. (continued)

ACMG/AMP criteria code	Original ACMG/AMP rule summary	Specification	Stand-alone	Very strong	Strong	Moderate	Supporting	Comments
PP1	Cosegregation with disease in multiple affected family members	Disease-specific, strength	NA	NA	Seven or more meioses observed within 1 family or across multiple families	Five or 6 meioses observed within 1 family or across multiple families	Three or 4 meioses observed within 1 family or across multiple families	(1) Affected individuals exhibit at least 1 of the <i>RUNX1</i> -specific phenotypic criteria. (2) Only genotype- and phenotype-positive individuals and obligate carriers are counted. (3) Demonstration of cosegregation in multiple families is not required because many <i>RUNX1</i> variants are unique and only occur in 1 family
PP2	Missense variant in a gene that has a low rate of BEN missense variation and where missense variants are a common mechanism of disease	NA	NA	NA	NA	NA	NA	Missense constraint z score for <i>RUNX1</i> is <3.09
PP3	Multiple lines of computational evidence support a deleterious effect on the gene or gene product	General recommendation	NA	NA	NA	NA	Per original ACMG/AMP guidelines	(1) PP3 should be applied for missense variants with a REVEL score >0.75. (2) PP3 should be applied for missense or synonymous variants if the variant alters the last 3 bases of an exon preceding a donor splice site or the first 3 bases of an exon following a splice acceptor site and the predicted decrease in the score of the canonical splice site (measured by both MES and SSF) is at least 75% regardless of the predicted creation/presence of a putative cryptic splice site. (3) PP3 should also be applied for intronic variants (in introns 4-8) located in reference to exons at positions +3 to +5 for splice donor sites or -3 to -5 for splice acceptor sites for which the predicted decrease in the score is at least 75% (measured by both MES and SSF) regardless of the predicted creation/presence of a putative cryptic splice site. (4) PP3 cannot be applied for canonical splice site variants
PP4	Patient's phenotype or family history is highly specific for a disease with a single genetic etiology	NA	NA	NA	NA	NA	NA	FPD/AML does not exhibit a highly specific phenotype, and there is substantial genetic heterogeneity

Evidence codes are noted in bold font.

AA, amino acid; co-IP, immunoprecipitation; EMSA, electrophoretic mobility shift assay; FRET, fluorescence resonance energy transfer; IF, immunofluorescence; LOF, loss-of-function; MES, MaxEntScan; NA, not applicable; NMD, nonsense-mediated decay; RHD, Runt homology domain; SSF, Splice Site Finder; WB, western blot; wt, wild type.

Table 1. (continued)

ACMG/AMP criteria code	Original ACMG/AMP rule summary	Specification	Stand-alone	Very strong	Strong	Moderate	Supporting	Comments
PP5	Reputable source recently reports variant as PATH, but the evidence is not available to the laboratory to perform an independent analysis	NA						According to SVI recommendations
BA1	Allele frequency is >5% in ESP, 1000G, or ExAC	Disease-specific	MAF ≥ 0.0015 (0.15%)	NA	NA	NA	NA	The variant is present in any general continental population dataset with a minimum number of 2000 alleles and variant present in ≥ 5 alleles
BS1	Allele frequency is greater than expected for disorder	Disease-specific	NA	NA	MAF between 0.00015 (0.015%) and 0.0015 (0.15%)	NA	NA	(1) The variant is present in any general continental population dataset with a minimum number of 2000 alleles and variant present in ≥ 5 alleles. (2) Variant can be classified as LBEN based on BS1 alone if there is no contradictory evidence supporting pathogenicity
BS2	Observed in a healthy adult individual for a recessive (homozygous), dominant (heterozygous), or X-linked (hemizygous) disorder, with full penetrance expected at an early age	NA						Patients with FPD/AML display incomplete penetrance, and the average age of onset of hematologic malignancies is 33 y
BS3	Well-established in vitro or in vivo functional studies show no damaging effect on protein function or splicing	Gene-specific, strength	NA	NA	(1) Transactivation assays exhibiting normal transactivation (80%-115% of wt); and (2) data from a secondary assay exhibiting normal function	NA	Transactivation assays exhibiting normal transactivation (80%-115% of wt)	See PS3 (1) and (2)
BS4	Lack of segregation in affected members of a family	General recommendation	NA	NA	Applied when seen in ≥ 2 informative meioses	NA	NA	This code should only be applied for genotype-positive, phenotype-negative (with sufficient laboratory evidence) family members
BP1	Missense variant in a gene for which primarily truncating variants are known to cause disease	NA						FDP/AML is caused by both PATH missense and truncating variants
BP2	Observed in trans with a PATH variant for a fully penetrant dominant gene/disorder or observed in cis with a PATH variant in any inheritance pattern	General recommendation	NA	NA	NA	NA	Per original ACMG/AMP guidelines	BP2 can also be applied if the variant is detected in a homozygous state
BP3	In-frame deletions/insertions in a repetitive region without a known function	NA						RUNX1 does not contain a repetitive region without known function

Evidence codes are noted in bold font.

AA, amino acid; co-IP, coimmunoprecipitation; EMSA, electrophoretic mobility shift assay; FRET, fluorescence resonance energy transfer; IF, immunofluorescence; LOF, loss-of-function; MES, MaxEntScan; NA, not applicable; NMD, nonsense-mediated decay; RHD, Runt homology domain; SSF, Splice Site Finder; WB, western blot; wt, wild type.

Table 1. (continued)

ACMG/AMP criteria code	Original ACMG/AMP rule summary	Specification	Stand-alone	Very strong	Strong	Moderate	Supporting	Comments
BP4	Multiple lines of computational evidence suggest no impact on gene or gene product	General recommendation	NA	NA	NA	NA	Per original ACMG/AMP guidelines	BP4 should be applied for missense variants if all of the following apply: (1) REVEL score <0.15; (2) SSF and MES predict either an increase in the canonical splice site score or a decrease in the canonical splice site score by no more than 10%; and (3) no putative cryptic splice sites are created. BP4 should also be applied for synonymous, intronic, and noncoding variants for which SSF and MES predict either an increase in the canonical splice site score or a decrease in the canonical splice site score by no more than 10% and no putative cryptic splice sites are created
BP5	Variant found in a case with an alternate molecular basis for disease	NA						In rare circumstances, a patient can carry 2 variants in genes predisposing to hematologic malignancies
BP6	Reputable source recently reports variants as BEN, but the evidence is not available to the laboratory to perform an independent evaluation	NA						According to SVI recommendations
BP7	A synonymous variant for which splicing prediction algorithms predict no impact to the splice consensus sequence, nor the creation of a new splice site, and the nucleotide is not highly conserved	General recommendation	NA	NA	NA	NA	Per original ACMG/AMP guidelines. BP7 cannot be applied in combination with PP3	Also applicable to intronic/noncoding variants at or beyond positions +7/-21 for which (1) SSF and MES predict either an increase in the canonical splice site score or a decrease in the canonical splice site score by no more than 10% and no putative cryptic splice sites are created; and (2) evolutionary conservation prediction algorithms predict the site as not conserved (eg, PhyloP score <0.1 or the variant is the reference nucleotide in 1 primate and/or 3 mammal species)

Evidence codes are noted in bold font.

AA, amino acid; co-IP, coimmunoprecipitation; EMSA, electrophoretic mobility shift assay; FRET, fluorescence resonance energy transfer; IF, immunofluorescence; LOF, loss-of-function; MES, MaxEntScan; NA, not applicable; NMD, nonsense-mediated decay; RHD, Runt homology domain; SSF, Splice Site Finder; WB, western blot; wt, wild type.

Table 2. FPD/AML phenotypic criteria

Feature	Details	Lifetime risk
Thrombocytopenia	Mild to moderate, normal platelet size and volume, absence of other causes for thrombocytopenia	In most patients
Platelet ultrastructural and/or functional defects	Includes platelet alpha or dense granule secretion defects and impaired platelet aggregation (particularly in response to collagen and epinephrine)	Unknown
Hematologic malignancy	Most commonly AML or MDS, less frequently T-ALL. There are rare case reports of patients with germline <i>RUNX1</i> mutations and mixed MPN/MDS such as CMML, as well as case reports of patients with B-cell ALL and hairy-cell leukemia	~44%

CMML, chronic myelomonocytic leukemia; MPN, myeloproliferative syndrome.

evidence can be applied. The MM-VCEP adopted the approach taken by various ClinGen expert panels,⁴⁵⁻⁴⁸ and supported by the SVI and others,⁴⁹ that additional meioses support higher levels of evidence. Thus, based on a calculated logarithm of the odds score thresholds of 0.9, 1.5, and 2.1, respectively, 3 or 4 meioses fulfill criteria for PP1, 5 or 6 meioses for PP1_moderate, and ≥ 7 meioses for PP1_strong. Of note, only individuals well documented as having an *RUNX1* phenotype (Table 2) and a positive genotype or obligate carriers are included when counting segregations. The phenotype of those individuals should be well described. We waived the ACMG/AMP recommendation for demonstrating cosegregation in >1 family, given that many *RUNX1* variants are unique to a single family¹³ and have not been reported in other unrelated families, which would severely affect the utility of segregation data. We acknowledge that by waiving this recommendation, there is a possibility of the identified variant being in a linkage disequilibrium with a truly causative variant.

Lack of segregation in affected family members (BS4) can be used as a BEN criterion when an *RUNX1* variant is present and nonsegregation with disease occurred in at least 2 or more informative meioses. BS4 should only be applied for genotype-positive, phenotype-negative family members, and there must be confidence that the family members do not meet any of our *RUNX1*-phenotypic criteria, taking into account ages of individuals.

De novo occurrence (PS2_moderate, PS2_supporting, PM6, and PM6_supporting)

De novo *RUNX1* variants are rare but have been reported in the literature.¹⁶⁻¹⁸ The 2 de novo criteria are applied when both maternity and paternity are confirmed (PS2) or assumed (PM6) and the variant has been assessed as de novo in a patient with the disease and no family history. The following specifications were added by our MM-VCEP: (1) no family history is defined by the absence of the FPD/AML-specific phenotype in first- and/or second-degree relatives; and (2) the proband must exhibit at least 1 phenotypic FPD/AML criterion (Table 2). PS2/PM6 were further specified by using the SVI recommendation of a point-based scoring system to determine the level of strength. The FPD/AML phenotype is not highly specific, and there is substantial genetic heterogeneity; the same phenotype can be caused by other underlying germline conditions such as *PATH* variants in *ANKRD26*⁵⁰ or *ETV6*.⁵¹ We thus concluded that due to the lack of a highly specific phenotype and the presence of genetic heterogeneity, the maximum allowable value is 1 point contributing

to the overall score. Due to this restriction, these 2 criteria do not have a strong or very strong level of evidence. PS2_moderate is reached with a score of 1 point (two or more proven de novo occurrences), and PS2_supporting is used when reaching a score of 0.5 point (one proven de novo occurrence). Likewise, PM6_moderate is met when 4 assumed de novo occurrences are present (score of 1), and PM6_supporting is applicable with 2 to 3 assumed de novo cases (score of 0.5). Combining these 2 criteria (eg, in the case of the same variant having both confirmed and assumed de novo evidence) is possible with the recognition that the maximum allowable value is still 1 point, which effectively leads to the application of 1 moderate or 2 supporting rules (supplemental Table 2).

Computational and predictive data (PVS1, PVS1_strong, PVS1_moderate, PS1, PS1_moderate, PM1, PM1_supporting, PM4, PM4_supporting, PM5_strong, PM5, PM5_supporting, PP3, BP4, and BP7)

RUNX1 germline variants have been well described as being dominant-negative, loss-of-function, or hypermorphic.^{13,31,52,53} Three major isoforms (A, B, and C) are expressed by the use of 2 promoters and alternative splicing (Figure 1B). Expression of the short human *RUNX1A* isoform has been shown to favor expansion of the hematopoietic stem cell pool, whereas expression of the full-length *RUNX1B* and *RUNX1C* isoforms, which only differ by 33 AAs at the N terminus of isoform C (exons 2-3 in NM_001754.4), function to promote hematopoietic differentiation.⁵⁴⁻⁶¹ The differential function and expression of these isoforms in hematopoietic tissue are not fully understood. The MM-VCEP recommends using *RUNX1* isoform C as the default transcript (NM_001754.4) because this is the isoform used for annotation by most clinical laboratories. The MM-VCEP decision tree for SNVs/indels (Figure 2; supplemental Table 3) and CNVs (supplemental Figure 1) refined the PVS1 criterion across all loss-of-function variant types previously reported for *RUNX1* (nonsense, frameshift, canonical splice site variants, and single- or multi-exon deletions) by using the SVI recommendations²⁷ and gene-specific adjustments. We recommend downgrading the strength level from very strong to strong for C-terminal variants that are not predicted to undergo nonsense-mediated decay but affect the transactivation domain, inhibitory domain, and/or the VWRPY motif (Figure 1).^{62,63} Nonsense-mediated decay is not predicted if the premature termination codon occurs in the 3'-most exon or within the 3'-most 50 nucleotides of the penultimate




Figure 1. Schematic of *RUNX1* exon distribution, protein isoforms, and functional domain structure with all 52 pilot variants and their final MM-VCEP classification. (A) Isoform C with RHD, transactivation domain (TAD), and the VWRPY motif and location of all 49 single-nucleotide pilot variants with their final MM-VCEP classification. PATH and LPATH variants are shown at the top, and VUS, LBEN, and BEN variants are shown at the bottom. The exonic distribution of isoform C is displayed below. (B) Schematic of *RUNX1* isoforms A, B, and C and their functional domains. Regions in gray are unique to 1 isoform. The 3 pilot CNVs are shown at the bottom, with the deletion of exons 2 and 3 exclusively affecting the N-terminal 33 AA of isoform C.

exon.^{64,65} Deletions of exon 2-3, presumably only affecting *RUNX1* isoform C, have been reported in 4 families (L.A.G., A.L.B., P.B., and D.P., written communication, 1 July 2019),⁶⁶ displaying a typical FPD/AML phenotype and segregation with disease. Although the functional effects of the exon 2-3 deletions on isoform C and potential effects on isoforms A and B require further investigation, we recommend applying PVS1_moderate according to the PVS1 CNV decision tree. The ClinGen CNV interpretation working group is currently developing a systematic framework for the clinical interpretation of CNVs, which will benefit the future curation of *RUNX1* CNVs.

A variant affecting the same AA residue as a previously established PATH variant can either lead to the same AA change (PS1) or

a different AA change (PM5). The MM-VCEP added the following recommendations for both rules: RNA data, or agreement in splicing predictors showing no splicing effects, which was defined as SSF and MES predicting either an increase in the canonical splice site score or a decrease in the canonical splice site score by no more than 10% and no putative cryptic splice sites are created. In addition, the previously established variant must be asserted PATH/LPATH based on MM-VCEP rules for *RUNX1* before this rule can be applied. A strength modification was established for PS1 (same AA as previously established PATH variant) and PS1_moderate (same AA as previously established LPATH variant). Likewise, PM5_strong is applied when 2 or more different PATH missense changes have been detected previously at the same AA




Figure 2. PVS1 decision tree for SNVs/indels. Application of different levels of strength for PVS1 depending on the prediction of nonsense-mediated decay (NMD), the location within a known critical protein domain, and the expression of alternative isoforms. The splicing effects table is given in supplemental Data.

residue, PM5 is used when a different PATH missense change has been seen previously at the same residue, and PM5_supporting is used when one missense change at the same residue has previously been determined to be LPATH.

For in silico evaluation of missense variants, the MM-VCEP recommends using REVEL, a meta-predictor that combines 13 individual tools with high sensitivity and specificity and has recently shown the highest performance compared with any individual tool or other ensemble methods.^{67,68} For splicing predictions, we recommend using the SSF and MES, both of which have been shown to predict splicing effects with high accuracy.⁶⁹⁻⁷¹ PP3, defined as multiple lines of computational evidence supporting a deleterious effect, can be applied for missense or synonymous variants if the variant alters the last 3 bases of an exon preceding a splice donor site or the first 3 bases of an exon following a splice acceptor site,⁶⁹ and if the predicted decrease in the score of the canonical splice site (measured by both MES and SSF) is at least 75% regardless of the predicted creation/presence of a putative cryptic splice site. PP3 should be applied for intronic variants (in introns 4-8) located in reference to exons at positions +3 to +5 for splice donor sites or -3 to -5 for splice acceptor sites^{69,72} for which the predicted decrease in the score of the canonical splice site is at least 75% (measured by both MES and SSF) regardless of the predicted creation/presence of a putative cryptic splice site. PP3 cannot be used for canonical splice site variants.

The BEN criterion BP4 should be applied for missense variants if all of the following criteria apply: the variant's REVEL score is <0.15, SSF and MES predict either an increase in the canonical splice site

score or a decrease in the canonical splice site score by no more than 10%, and no putative cryptic splice sites are created. BP4 should also be applied for synonymous, intronic, and noncoding variants for which SSF and MES predict either an increase in the canonical splice site score or a decrease in the canonical splice site score by no more than 10%, and no putative cryptic splice sites are created.

The original PM1 code can be applied for variants affecting mutational hotspots and/or functional domains without BEN variation. The RHD, spanning from AA 77-204, has been established as a highly conserved DNA-binding domain without any BEN variation in ClinVar. Thirteen somatic and/or germline mutational hotspots within the RHD have been identified: R107, K110, A134, R162, R166, S167, R169, G170, K194, T196, D198, R210, and R204.^{12,20,73,74} The MM-VCEP recommends using PM1 for variants affecting these 13 AA residues. For variants in other parts of the RHD for which germline variants have been previously reported (AA 105-204), a reduced-strength level (PM1_supporting) is recommended. For other residues within the RHD (AA 77-104), no germline *RUNX1* PATH variants have been reported to date. In the future, the AA range under PM1_supporting may be expanded to other parts of the protein if more evidence emerges. Analogous to PM1, PM4 (protein length changes due to in-frame deletions/insertions in a nonrepeat region or stop-loss variants) is applied to in-frame deletions/insertions affecting the same 13 AA residues (as listed earlier) and, likewise, PM4_supporting can be used for in-frame deletions/insertions affecting at least one of the other parts of the RHD in which germline variants have been previously reported (AA 105-204).

The MM-VCEP agreed to extend BP7 (synonymous variant with no splicing effect and position is not highly conserved) to apply to intronic/noncoding variants at or beyond positions +7/-21 for

which SSF and MES predict either an increase in the canonical splice site score or a decrease in the canonical splice site score by no more than 10%, no putative cryptic splice sites are created, and the position is not conserved (eg, PhyloP score $<0.1^{75}$) or the variant is the reference nucleotide in 1 primate and/or 3 mammal species.⁷⁶

Functional data (PS3, PS3_moderate, PS3_supporting, BS3, and BS3_supporting)

The evolutionarily conserved 128 AA RHD, present in most of the RUNX1 isoforms (Figure 1B), is involved in DNA binding and heterodimerization with core binding factor (CBF) β . Heterodimerization of RUNX1 with CBF β promotes DNA binding by stabilizing the interaction of the complex with the DNA. RUNX1 regulates the activity of several important hematopoietic genes, such as the granulocyte-macrophage colony-stimulating factor,^{77,78} T-cell receptor,^{79,80} myeloperoxidase,^{81,82} and neutrophil elastase,⁸² by binding to a core sequence (TGTGGT) found in their promoters or enhancers.

Transactivation assays showing altered transactivation compared with wild type are often performed as functional studies to evaluate the pathogenicity of a RUNX1 variant. Promoter sequences of *M-CSFR*, *PF4*, *C-FMS*, and *GZMB*, containing consensus RUNX1 binding sites TGTGGT have been used for this purpose.^{31,83-87} Data from secondary assays are frequently used to evaluate an altered function of mutant RUNX1. Electrophoretic mobility shift assays^{31,87-90} and yeast hybrid assays^{88,89} are performed to show decreased DNA-binding affinity, and coimmunoprecipitation assays,^{85,87,90} fluorescence resonance energy transfer assays,⁸⁸ and affinity assays³¹ can illustrate the diminished heterodimerization ability of mutant RUNX1 with CBF β . Abnormal cellular localization of mutant RUNX1 can be shown by immunofluorescence^{31,53,83} and cell fractionation with western blot.^{85,90} Sorted primary hematopoietic stem and progenitor cells can be used to show reduced colony-forming potential,^{53,91} and xenotransplantation experiments may reveal abnormal function of mutant RUNX1 in vivo.⁵³

The MM-VCEP defined the strong PATH code PS3 as the combination of reduced transactivation ($<20\%$ of wild type and/or reduced to levels similar to well-established PATH variants such as R201Q or R166Q) and data from a secondary assay that show altered function of mutant RUNX1. The transactivation assay should include wild-type and known PATH controls as well as coexpression with CBF β . PS3 can also be applied for evidence of very low or abnormal messenger RNA (mRNA)/protein expression of the variant allele as a functional consequence of a null variant or incorrect mRNA/protein products. The MM-VCEP further stipulates that PS3 cannot be applied if the variant meets PVS1. If the variant meets PVS1_strong and PS3, we recommend applying either PVS1_strong and PS3_moderate or upgrading PVS1_strong to PVS1 without applying PS3. PS3_moderate is applied when data from transactivation assays exhibit reduced transactivation ($<20\%$ of wild type and/or reduced to levels similar to well-established PATH variants such as R201Q or R166Q) or 2 or more secondary assays show altered function. PS3_supporting can be applied for transactivation assays exhibiting enhanced transactivation ($>115\%$ of wild type), as has been reported previously for the hypermorphic RUNX1 mutant, S388X.⁵²

Likewise, the BS3 requirements (functional studies show no damaging effect on protein function) are a normal transactivation

(80%-115% of wild type) and data from a secondary assay that exhibit normal function. BS3_supporting can be applied when there is evidence of normal transactivation (80%-115% of wild type); data from secondary assays are not required.

Rules deemed not applicable

Four rules of the PATH framework (PM3, PP2, PP4, and PP5) and 5 rules of the BEN framework (BS2, BP1, BP3, BP5, and BP6) were deemed not applicable. The reasoning behind the decision for each code is briefly explained here.

Because the FPD/AML phenotype is associated with autosomal dominant transmission, PM3 (detected in trans with a PATH variant in a recessive gene) cannot be applied for FPD/AML.

The recommended cutoff for PP2 (missense variant in a gene with low rate of missense variants) is a constraint z score ≥ 3.09 ,²⁸ which was not met by RUNX1.

The phenotype observed in FPD/AML is rather nonspecific and can be caused by a number of other inherited predisposition syndromes, somatic variants, or environmental factors; this scenario makes the original ACMG/AMP rule PP4 for a highly specific phenotype not applicable to RUNX1.

Incomplete penetrance, an average age of onset of 33 years for hematologic malignancies,¹⁹⁻²¹ and the lack of sufficient clinical data to exclude an RUNX1-related phenotype render BS2 (observed in a healthy individual with full penetrance at an early age) not applicable.

Both missense and truncating variants have been described as causative in FPD/AML, making BP1 (missense variant in a gene with primarily truncating variants) not applicable. Similarly, BP3 is not applicable, as RUNX1 lacks repetitive regions of unknown function.


BP5 can be applied when the variant is found in a case with an alternate molecular basis for disease. The MM-VCEP concluded that this rule is not applicable because in rare circumstances, a patient can carry variants in 2 genes predisposing to hematologic malignancies, as has been described in case reports. In addition, variants in other genes presenting as low-penetrance risk factors, modifier genes, and/or somatic mutations in hematopoietic stem and progenitor cells may contribute to the clinical presentation and complicate the search for the causative variant.^{92,93}

Following recommendations from the SVI, the MM-VCEP agreed not to use the 2 variant classifications from reputable source evidence codes (PP5 and BP6) based on the published rationale.²⁸

Performance of the MM-VCEP specifications in pilot variant classification

For pilot testing, 52 variants with a broad spectrum of ClinVar assertions (12 BEN/LBEN, 14 VUS, 20 PATH/LPATH, 4 CONF, and 2 variants with no ClinVar assertions) were selected. The MM-VCEP applied the RUNX1-modified ACMG/AMP criteria to all pilot variants. During testing, experts were able to provide feedback on the usability of the evidence codes, comment on the weight of certain lines of evidence, and suggest further modifications of the rule. A list of all pilot variants, the variant classification of the ClinVar submitters, and the classifications made by our MM-VCEP are presented in supplemental Table 4. Figure 3 compares the original ClinVar classifications vs our MM-VCEP classifications grouped

Figure 3. Comparison of ClinVar and MM-VCEP classifications. Fifty previously asserted and ClinVar-deposited *RUNX1* variants are shown on the x-axis. Final MM-VCEP classifications are color-coded (see legend on the right). ClinVar variants with previous LPATH, CONF, and VUS assertions were most often reclassified by using MM-VCEP-specified rules for *RUNX1*.



according to PATH/LPATH, BEN/LBEN, VUS, and CONF variants. Of the 14 VUS, 2 were upgraded into the LPATH category. Of the 4 CONF variants, 1 was upgraded to PATH, and 3 were downgraded to LBEN. Two of 18 variants previously listed as PATH/LPATH in ClinVar were downgraded to VUS after applying the *RUNX1*-specific codes. MM-VCEP members with knowledge of the criteria applied by the ClinVar submitters were able to corroborate the VUS classifications. A detailed schematic of the *RUNX1* gene and the newly classified pilot variants is shown in Figure 1A. Overall, applying the *RUNX1* specifications to the VUS/CONF variants resulted in a reduction in VUS/CONF classifications of 33%. All of the 12 variants that were submitted in ClinVar as BEN/LBEN remained in this category, with most LBEN variants being downgraded to BEN and only 2 remaining LBEN. An overview of the frequency of PATH and BEN evidence codes applied is given in supplemental Figure 2. The test set received a final concordance of 92% with consensus ClinVar classifications (90% for the PATH/LPATH test set, 86% with the VUS test set, and 100% for the BEN/LBEN test set).

Discussion

RUNX1 is commonly mutated in hematologic malignancies with high rates of somatic variants in MDS/AML.^{73,94} Tumor-based next-generation sequencing panels covering *RUNX1* among other genes are implicated in the molecular diagnostic process of MDS/AML in most treatment centers. Some of these somatic *RUNX1* variants are subsequently determined to be germline.⁹⁵ In addition, recent achievements such as the inclusion of inherited hematologic malignancies into the revised World Health Organization classification of myeloid neoplasms and acute leukemia¹⁵ and a more standardized evaluation of family history have raised awareness of these syndromes among physicians. This awareness will increase the identification of patients with FPD/AML. Accurate *RUNX1* variant curation is fundamental for the appropriate clinical care of these patients, especially when considering a related donor for hematopoietic stem cell transplantation. In addition, FPD/AML with thrombocytopenia may be misdiagnosed as immune thrombocytopenic purpura, and the correlating dysmegakaryopoiesis in the bone marrow can be mistaken as an early-stage MDS, underscoring the importance of adequate *RUNX1* variant curation.^{96,97}

Our curation of pilot variants showed the impact of our proposed rules on improving variant classification, resulting in a reduction of VUS/CONF variants by 33%. Further use of these rules should continue to reduce the number of VUS and lead to fewer number of variants with VUS/CONF assertion within ClinVar. Being able to reclassify a variant from VUS/CONF assertions has a significant impact on patient care as it provides patients and physicians with the definitive data to guide treatment decisions, including donor selection among matched relatives. As we implement these *RUNX1*-specific rules, the variant annotation in ClinVar will contain a link to the specific version of the MM-VCEP *RUNX1* evidence rules, a summary of the specific evidence codes used for that variant, and a link to the ClinGen evidence repository where all the evidence evaluated for that variant is found. Given these detailed expert-reviewed curations, MM-VCEP-curated variants will be submitted under a “3-star expert panel reviewed” FDA-recognized designation.

We expect that our *RUNX1*-specific rules will require further updating as additional data become available, or at a minimum every 2 years, and will address improved computational modeling, functional assays, and larger and more ethnically diverse population databases. Per ClinGen policy, *RUNX1* VUS and LPATH variants will be reassessed by the expert panel every 2 years, and other variants may be re-curated if discrepancies in the variant classification or new evidence emerge over time. At any time, a link to the most up-to-date recommendations of *RUNX1* evidence codes can be found on the MM-VCEP home page (<https://www.clinicalgenome.org/affiliation/50034>). Furthermore, ongoing general refinements to the ACMG/AMP guidelines made by the ClinGen SVI will need to be addressed, particularly for the curation of intragenic *RUNX1* deletions and consensus rules for evaluation of splicing predictions. The next step of the MM-VCEP will be the curation of all current ClinVar-deposited *RUNX1* variants. Further work will extend this study to other genes causing inherited hematologic malignancies.

Acknowledgments

The VCEP thanks the ClinGen SVI Working Group as well as the Executive Committee of the Hereditary Cancer Clinical Domain Working Group.

Results provided in this publication were generated by the American Society of Hematology in collaboration with Baylor College of Medicine and the University of North Carolina, NIH-funded Clinical Genome Resource grant award recipients. The NIH, National Human Genome Research Institute supported this work through U41HG009649 (X.L. and S.E.P.) and U41HG009650 (S.M. and J.E.R.); and the 2018 NIH/National Cancer Institute Leukemia SPORE DRP award (P50CA100632-16, project 00007529) (C.D.D.).

Authorship

Contribution: All of the authors participated in the construction and pilot testing of the *RUNX1* curation rules and edited the manuscript; and X.L., S.F., S.M., D.W., and L.A.G. participated in the majority of the manuscript writing.

Conflict-of-interest disclosure: S.E.P. is a member of the scientific advisory panel of Baylor Genetics Laboratories. L.A.G. is a member of the scientific advisory board for Invitae, Inc., and receives royalties from UpToDate, Inc. L.Z. received honoraria from Future Technology Research, LLC, Roche Diagnostics Asia Pacific,

BGI, and Illumina. A family member of L.Z. has a leadership position and ownership interest in the Shanghai Genome Center. The remaining authors declare no competing financial interests.

ORCID profiles: S.M., 0000-0002-2796-282X; C.C.P., 0000-0001-8774-0180; A.L.B., 0000-0002-9023-0138; C.K., 0000-0001-6043-2065; M.L., 0000-0002-5788-3962; J.E.R., 0000-0002-7437-8060; C.D.D., 0000-0001-9003-0390; A.A.B., 0000-0003-1864-8502; N.M., 0000-0003-1465-2934; K.E.N., 0000-0002-5581-6555; J.S., 0000-0002-8734-5356; A.R., 0000-0002-7368-2247; M.J.R., 0000-0003-4860-261X; L.Z., 0000-0003-4517-0751; N.A.S., 0000-0002-1893-582X; S.E.P., 0000-0002-9626-0936; D.W., 0000-0001-7729-5730; L.A.G., 0000-0003-1914-9158.

Correspondence: David Wu, University of Washington, 825 Eastlake Ave E, G7800, Seattle, WA 98109; e-mail: dwu2@uw.edu; and Lucy A. Godley, The University of Chicago, 5841 S Maryland Ave, MC 2115, Chicago, IL 60637; e-mail: lgodley@medicine.bsd.uchicago.edu.

References

1. Richards S, Aziz N, Bale S, et al; ACMG Laboratory Quality Assurance Committee. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424.
2. Pepin MG, Murray ML, Bailey S, Leistriz-Kessler D, Schwarze U, Byers PH. The challenge of comprehensive and consistent sequence variant interpretation between clinical laboratories. *Genet Med*. 2016;18(1):20-24.
3. Amendola LM, Jarvik GP, Leo MC, et al. Performance of ACMG-AMP variant-interpretation guidelines among nine laboratories in the Clinical Sequencing Exploratory Research Consortium [published correction appears in *Am J Hum Genet*. 2016;99(1):247]. *Am J Hum Genet*. 2016;98(6):1067-1076.
4. Harrison SM, Dolinsky JS, Knight Johnson AE, et al. Clinical laboratories collaborate to resolve differences in variant interpretations submitted to ClinVar. *Genet Med*. 2017;19(10):1096-1104.
5. MacArthur DG, Manolio TA, Dimmock DP, et al. Guidelines for investigating causality of sequence variants in human disease. *Nature*. 2014;508(7497):469-476.
6. Whiffin N, Minikel E, Walsh R, et al. Using high-resolution variant frequencies to empower clinical genome interpretation. *Genet Med*. 2017;19(10):1151-1158.
7. Song W, Gardner SA, Hovhannisyan H, et al. Exploring the landscape of pathogenic genetic variation in the ExAC population database: insights of relevance to variant classification. *Genet Med*. 2016;18(8):850-854.
8. Harrison SM, Dolinsky JS, Chen W, et al; ClinGen Sequence Variant Inter-Laboratory Discrepancy Resolution Working Group. Scaling resolution of variant classification differences in ClinVar between 41 clinical laboratories through an outlier approach. *Hum Mutat*. 2018;39(11):1641-1649.
9. Rivera-Muñoz EA, Milko LV, Harrison SM, et al. ClinGen Variant Curation Expert Panel experiences and standardized processes for disease and gene-level specification of the ACMG/AMP guidelines for sequence variant interpretation. *Hum Mutat*. 2018;39(11):1614-1622.
10. Rehm HL, Berg JS, Brooks LD, et al; ClinGen. ClinGen—the Clinical Genome Resource. *N Engl J Med*. 2015;372(23):2235-2242.
11. Landrum MJ, Lee JM, Benson M, et al. ClinVar: public archive of interpretations of clinically relevant variants. *Nucleic Acids Res*. 2016;44(D1):D862-D868.
12. Schlegelberger B, Heller PG. *RUNX1* deficiency (familial platelet disorder with predisposition to myeloid leukemia, FPDMM). *Semin Hematol*. 2017;54(2):75-80.
13. Sood R, Kamikubo Y, Liu P. Role of *RUNX1* in hematological malignancies. *Blood*. 2017;129(15):2070-2082.
14. Song WJ, Sullivan MG, Legare RD, et al. Haploinsufficiency of *CBFA2* causes familial thrombocytopenia with propensity to develop acute myelogenous leukaemia. *Nat Genet*. 1999;23(2):166-175.
15. Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016;127(20):2391-2405.
16. Béri-Dexheimer M, Latger-Cannard V, Philippe C, et al. Clinical phenotype of germline *RUNX1* haploinsufficiency: from point mutations to large genomic deletions. *Eur J Hum Genet*. 2008;16(8):1014-1018.
17. Schmit JM, Turner DJ, Hromas RA, et al. Two novel *RUNX1* mutations in a patient with congenital thrombocytopenia that evolved into a high grade myelodysplastic syndrome. *Leuk Res Rep*. 2015;4(1):24-27.

18. Ouchi-Uchiyama M, Sasahara Y, Kikuchi A, et al. Analyses of genetic and clinical parameters for screening patients with inherited thrombocytopenia with small or normal-sized platelets. *Pediatr Blood Cancer*. 2015;62(12):2082-2088.
19. Feurstein S, Drazer MW, Godley LA. Genetic predisposition to leukemia and other hematologic malignancies. *Semin Oncol*. 2016;43(5):598-608.
20. Godley LA. Inherited predisposition to acute myeloid leukemia. *Semin Hematol*. 2014;51(4):306-321.
21. West AH, Godley LA, Churpek JE. Familial myelodysplastic syndrome/acute leukemia syndromes: a review and utility for translational investigations. *Ann N Y Acad Sci*. 2014;1310(1):111-118.
22. Buijs A, Poddighe P, van Wijk R, et al. A novel CBFA2 single-nucleotide mutation in familial platelet disorder with propensity to develop myeloid malignancies. *Blood*. 2001;98(9):2856-2858.
23. Kiritto K, Sakoe K, Shinoda D, Takiyama Y, Kaushansky K, Komatsu N. A novel RUNX1 mutation in familial platelet disorder with propensity to develop myeloid malignancies. *Haematologica*. 2008;93(1):155-156.
24. Hamilton KV, Maese L, Marron JM, Pulsipher MA, Porter CC, Nichols KE. Stopping leukemia in its tracks: should preemptive hematopoietic stem-cell transplantation be offered to patients at increased genetic risk for acute myeloid leukemia? *J Clin Oncol*. 2019;37(24):2098-2104.
25. Churpek JE, Godley LA; University of Chicago Hematopoietic Malignancies Cancer Risk Team. How I diagnose and manage individuals at risk for inherited myeloid malignancies. *Blood*. 2016;128(14):1800-1813.
26. Owen CJ, Toze CL, Koochin A, et al. Five new pedigrees with inherited RUNX1 mutations causing familial platelet disorder with propensity to myeloid malignancy. *Blood*. 2008;112(12):4639-4645.
27. Abou Tayoun AN, Pesaran T, DiStefano MT, et al; ClinGen Sequence Variant Interpretation Working Group (ClinGen SVI). Recommendations for interpreting the loss of function PVS1 ACMG/AMP variant criterion. *Hum Mutat*. 2018;39(11):1517-1524.
28. Biesecker LG, Harrison SM; ClinGen Sequence Variant Interpretation Working Group. The ACMG/AMP reputable source criteria for the interpretation of sequence variants. *Genet Med*. 2018;20(12):1687-1688.
29. Tavtigian SV, Greenblatt MS, Harrison SM, et al; ClinGen Sequence Variant Interpretation Working Group (ClinGen SVI). Modeling the ACMG/AMP variant classification guidelines as a Bayesian classification framework. *Genet Med*. 2018;20(9):1054-1060.
30. Latger-Cannard V, Philippe C, Bouquet A, et al. Haematological spectrum and genotype-phenotype correlations in nine unrelated families with RUNX1 mutations from the French network on inherited platelet disorders. *Orphanet J Rare Dis*. 2016;11(1):49.
31. Michaud J, Wu F, Osato M, et al. In vitro analyses of known and novel RUNX1/AML1 mutations in dominant familial platelet disorder with predisposition to acute myelogenous leukemia: implications for mechanisms of pathogenesis. *Blood*. 2002;99(4):1364-1372.
32. Stockley J, Morgan NV, Bem D, et al; UK Genotyping and Phenotyping of Platelets Study Group. Enrichment of FLI1 and RUNX1 mutations in families with excessive bleeding and platelet dense granule secretion defects. *Blood*. 2013;122(25):4090-4093.
33. Johnson B, Lowe GC, Futterer J, et al; UK GAPP Study Group. Whole exome sequencing identifies genetic variants in inherited thrombocytopenia with secondary qualitative function defects. *Haematologica*. 2016;101(10):1170-1179.
34. Walker LC, Stevens J, Campbell H, et al. A novel inherited mutation of the transcription factor RUNX1 causes thrombocytopenia and may predispose to acute myeloid leukaemia. *Br J Haematol*. 2002;117(4):878-881.
35. Jalagadugula G, Mao G, Kaur G, Goldfinger LE, Dhanasekaran DN, Rao AK. Regulation of platelet myosin light chain (MYL9) by RUNX1: implications for thrombocytopenia and platelet dysfunction in RUNX1 haploinsufficiency. *Blood*. 2010;116(26):6037-6045.
36. Nishimoto N, Imai Y, Ueda K, et al. T cell acute lymphoblastic leukemia arising from familial platelet disorder. *Int J Hematol*. 2010;92(1):194-197.
37. Preudhomme C, Renneville A, Bourdon V, et al. High frequency of RUNX1 biallelic alteration in acute myeloid leukemia secondary to familial platelet disorder. *Blood*. 2009;113(22):5583-5587.
38. Shiba N, Hasegawa D, Park MJ, et al. CBL mutation in chronic myelomonocytic leukemia secondary to familial platelet disorder with propensity to develop acute myeloid leukemia (FPD/AML). *Blood*. 2012;119(11):2612-2614.
39. Linden T, Schnittger S, Groll AH, Juergens H, Rossig C. Childhood B-cell precursor acute lymphoblastic leukaemia in a patient with familial thrombocytopenia and RUNX1 mutation. *Br J Haematol*. 2010;151(5):528-530.
40. Toya T, Yoshimi A, Morioka T, et al. Development of hairy cell leukemia in familial platelet disorder with predisposition to acute myeloid leukemia. *Platelets*. 2014;25(4):300-302.
41. Horowitz GL, Atlaie S, Boyd JC. Defining, establishing, and verifying reference intervals in the clinical laboratory; approved guideline. Wayne, PA: Clinical and Laboratory Standards Institute; 2010.C28-A3c.
42. Ghosh R, Harrison SM, Rehm HL, Plon SE, Biesecker LG; ClinGen Sequence Variant Interpretation Working Group. Updated recommendation for the benign stand-alone ACMG/AMP criterion. *Hum Mutat*. 2018;39(11):1525-1530.
43. Wang Q, Stacy T, Binder M, Marin-Padilla M, Sharpe AH, Speck NA. Disruption of the Cbfa2 gene causes necrosis and hemorrhaging in the central nervous system and blocks definitive hematopoiesis. *Proc Natl Acad Sci USA*. 1996;93(8):3444-3449.
44. Okuda T, van Deursen J, Hiebert SW, Grosveld G, Downing JR. AML1, the target of multiple chromosomal translocations in human leukemia, is essential for normal fetal liver hematopoiesis. *Cell*. 1996;84(2):321-330.
45. Mester JL, Ghosh R, Pesaran T, et al. Gene-specific criteria for PTEN variant curation: recommendations from the ClinGen PTEN Expert Panel. *Hum Mutat*. 2018;39(11):1581-1592.
46. Kelly MA, Caleshu C, Morales A, et al. Adaptation and validation of the ACMG/AMP variant classification framework for MYH7-associated inherited cardiomyopathies: recommendations by ClinGen's Inherited Cardiomyopathy Expert Panel. *Genet Med*. 2018;20(3):351-359.

47. Oza AM, DiStefano MT, Hemphill SE, et al; ClinGen Hearing Loss Clinical Domain Working Group. Expert specification of the ACMG/AMP variant interpretation guidelines for genetic hearing loss. *Hum Mutat.* 2018;39(11):1593-1613.
48. Gelb BD, Cavé H, Dillon MW, et al; ClinGen RASopathy Working Group. ClinGen's RASopathy Expert Panel consensus methods for variant interpretation. *Genet Med.* 2018;20(11):1334-1345.
49. Jarvik GP, Browning BL. Consideration of cosegregation in the pathogenicity classification of genomic variants. *Am J Hum Genet.* 2016;98(6):1077-1081.
50. Pippucci T, Savoia A, Perrotta S, et al. Mutations in the 5' UTR of ANKRD26, the ankirin repeat domain 26 gene, cause an autosomal-dominant form of inherited thrombocytopenia, THC2. *Am J Hum Genet.* 2011;88(1):115-120.
51. Zhang MY, Churpek JE, Keel SB, et al. Germline ETV6 mutations in familial thrombocytopenia and hematologic malignancy. *Nat Genet.* 2015;47(2):180-185.
52. Churpek JE, Garcia JS, Madzo J, Jackson SA, Onel K, Godley LA. Identification and molecular characterization of a novel 3' mutation in RUNX1 in a family with familial platelet disorder. *Leuk Lymphoma.* 2010;51(10):1931-1935.
53. Bluteau D, Gilles L, Hilpert M, et al. Down-regulation of the RUNX1-target gene NR4A3 contributes to hematopoiesis deregulation in familial platelet disorder/acute myelogenous leukemia. *Blood.* 2011;118(24):6310-6320.
54. Challen GA, Goodell MA. Runx1 isoforms show differential expression patterns during hematopoietic development but have similar functional effects in adult hematopoietic stem cells. *Exp Hematol.* 2010;38(5):403-416.
55. Komeno Y, Yan M, Matsuura S, et al. Runx1 exon 6-related alternative splicing isoforms differentially regulate hematopoiesis in mice. *Blood.* 2014;123(24):3760-3769.
56. Brady G, Elgueta Karstegl C, Farrell PJ. Novel function of the unique N-terminal region of RUNX1c in B cell growth regulation. *Nucleic Acids Res.* 2013;41(3):1555-1568.
57. Lacaud G, Gore L, Kennedy M, et al. Runx1 is essential for hematopoietic commitment at the hemangioblast stage of development in vitro. *Blood.* 2002;100(2):458-466.
58. Navarro-Montero O, Ayllon V, Lamolda M, et al. RUNX1c regulates hematopoietic differentiation of human pluripotent stem cells possibly in cooperation with proinflammatory signaling. *Stem Cells.* 2017;35(11):2253-2266.
59. Lorsbach RB, Moore J, Ang SO, Sun W, Lenny N, Downing JR. Role of RUNX1 in adult hematopoiesis: analysis of RUNX1-IRES-GFP knock-in mice reveals differential lineage expression. *Blood.* 2004;103(7):2522-2529.
60. Tsuzuki S, Hong D, Gupta R, Matsuo K, Seto M, Enver T. Isoform-specific potentiation of stem and progenitor cell engraftment by AML1/RUNX1. *PLoS Med.* 2007;4(5):e172.
61. Tsuzuki S, Seto M. Expansion of functionally defined mouse hematopoietic stem and progenitor cells by a short isoform of RUNX1/AML1. *Blood.* 2012;119(3):727-735.
62. Liu H, Carlsson L, Grundström T. Identification of an N-terminal transactivation domain of Runx1 that separates molecular function from global differentiation function. *J Biol Chem.* 2006;281(35):25659-25669.
63. Nishimura M, Fukushima-Nakase Y, Fujita Y, et al. VWRPY motif-dependent and -independent roles of AML1/Runx1 transcription factor in murine hematopoietic development. *Blood.* 2004;103(2):562-570.
64. Chang YF, Imam JS, Wilkinson MF. The nonsense-mediated decay RNA surveillance pathway. *Annu Rev Biochem.* 2007;76(1):51-74.
65. Lewis BP, Green RE, Brenner SE. Evidence for the widespread coupling of alternative splicing and nonsense-mediated mRNA decay in humans. *Proc Natl Acad Sci U S A.* 2003;100(1):189-192.
66. Cavalcante de Andrade Silva M, Krepischi ACV, Kulikowski LD, et al. Deletion of RUNX1 exons 1 and 2 associated with familial platelet disorder with propensity to acute myeloid leukemia. *Cancer Genet.* 2018;222-223:32-37.
67. Ghosh R, Oak N, Plon SE. Evaluation of in silico algorithms for use with ACMG/AMP clinical variant interpretation guidelines. *Genome Biol.* 2017;18(1):225.
68. Ioannidis NM, Rothstein JH, Pejaver V, et al. REVEL: an ensemble method for predicting the pathogenicity of rare missense variants. *Am J Hum Genet.* 2016;99(4):877-885.
69. Houdayer C, Caux-Moncoutier V, Krieger S, et al. Guidelines for splicing analysis in molecular diagnosis derived from a set of 327 combined in silico/ in vitro studies on BRCA1 and BRCA2 variants. *Hum Mutat.* 2012;33(8):1228-1238.
70. Yeo G, Burge CB. Maximum entropy modeling of short sequence motifs with applications to RNA splicing signals. *J Comput Biol.* 2004;11(2-3):377-394.
71. Desmet F-O, Hamroun D, Lalonde M, Collod-Bérout G, Claustres M, Bérout C. Human splicing finder: an online bioinformatics tool to predict splicing signals. *Nucleic Acids Res.* 2009;37(9):e67.
72. Tang R, Prosser DO, Love DR. Evaluation of bioinformatic programmes for the analysis of variants within splice site consensus regions. *Adv Bioinforma.* 2016;2016:5614058.
73. Schnittger S, Dicker F, Kern W, et al. RUNX1 mutations are frequent in de novo AML with noncomplex karyotype and confer an unfavorable prognosis. *Blood.* 2011;117(8):2348-2357.
74. Tang JL, Hou HA, Chen CY, et al. AML1/RUNX1 mutations in 470 adult patients with de novo acute myeloid leukemia: prognostic implication and interaction with other gene alterations. *Blood.* 2009;114(26):5352-5361.

75. Pollard KS, Hubisz MJ, Rosenbloom KR, Siepel A. Detection of nonneutral substitution rates on mammalian phylogenies. *Genome Res.* 2010;20(1): 110-121.
76. Lee K, Krempely K, Roberts ME, et al. Specifications of the ACMG/AMP variant curation guidelines for the analysis of germline CDH1 sequence variants. *Hum Mutat.* 2018;39(11):1553-1568.
77. Cockerill PN, Osborne CS, Bert AG, Grotto RJ. Regulation of GM-CSF gene transcription by core-binding factor. *Cell Growth Differ.* 1996;7(7): 917-922.
78. Li X, Vradii D, Gutierrez S, et al. Subnuclear targeting of Runx1 is required for synergistic activation of the myeloid specific M-CSF receptor promoter by PU.1. *J Cell Biochem.* 2005;96(4):795-809.
79. Halle JP, Haus-Seuffert P, Woltering C, Stelzer G, Meisterernst M. A conserved tissue-specific structure at a human T-cell receptor beta-chain core promoter. *Mol Cell Biol.* 1997;17(8):4220-4229.
80. Redondo JM, Pfohl JL, Hernandez-Munain C, Wang S, Speck NA, Krangel MS. Indistinguishable nuclear factor binding to functional core sites of the T-cell receptor delta and murine leukemia virus enhancers. *Mol Cell Biol.* 1992;12(11):4817-4823.
81. Austin GE, Zhao WG, Regmi A, Lu JP, Braun J. Identification of an upstream enhancer containing an AML1 site in the human myeloperoxidase (MPO) gene. *Leuk Res.* 1998;22(11):1037-1048.
82. Nuchprayoon I, Meyers S, Scott LM, Suzow J, Hiebert S, Friedman AD. PEBP2/CBF, the murine homolog of the human myeloid AML1 and PEBP2 beta/CBF beta proto-oncogenes, regulates the murine myeloperoxidase and neutrophil elastase genes in immature myeloid cells. *Mol Cell Biol.* 1994;14(8): 5558-5568.
83. Osato M, Asou N, Abdalla E, et al. Biallelic and heterozygous point mutations in the runt domain of the AML1/PEBP2alphaB gene associated with myeloblastic leukemias. *Blood.* 1999;93(6):1817-1824.
84. Glembotsky AC, Bluteau D, Espasandin YR, et al. Mechanisms underlying platelet function defect in a pedigree with familial platelet disorder with a predisposition to acute myelogenous leukemia: potential role for candidate RUNX1 targets. *J Thromb Haemost.* 2014;12(5):761-772.
85. Zhao LJ, Wang YY, Li G, et al. Functional features of RUNX1 mutants in acute transformation of chronic myeloid leukemia and their contribution to inducing murine full-blown leukemia. *Blood.* 2012;119(12):2873-2882.
86. Koh CP, Wang CQ, Ng CEL, et al. RUNX1 meets MLL: epigenetic regulation of hematopoiesis by two leukemia genes. *Leukemia.* 2013;27(9): 1793-1802.
87. Tsai SC, Shih LY, Liang ST, et al. Biological activities of RUNX1 mutants predict secondary acute leukemia transformation from chronic myelomonocytic leukemia and myelodysplastic syndromes. *Clin Cancer Res.* 2015;21(15):3541-3551.
88. Matheny CJ, Speck ME, Cushing PR, et al. Disease mutations in RUNX1 and RUNX2 create nonfunctional, dominant-negative, or hypomorphic alleles. *EMBO J.* 2007;26(4):1163-1175.
89. Li Z, Yan J, Matheny CJ, et al. Energetic contribution of residues in the Runx1 Runt domain to DNA binding. *J Biol Chem.* 2003;278(35):33088-33096.
90. Okada Y, Watanabe M, Nakai T, et al. RUNX1, but not its familial platelet disorder mutants, synergistically activates PF4 gene expression in combination with ETS family proteins. *J Thromb Haemost.* 2013;11(9):1742-1750.
91. Antony-Debré I, Manchev VT, Balayn N, et al. Level of RUNX1 activity is critical for leukemic predisposition but not for thrombocytopenia. *Blood.* 2015; 125(6):930-940.
92. Riordan JD, Nadeau JH. From peas to disease: modifier genes, network resilience, and the genetics of health. *Am J Hum Genet.* 2017;101(2):177-191.
93. Senol-Cosar O, Schmidt RJ, Qian E, et al. Considerations for clinical curation, classification, and reporting of low-penetrance and low effect size variants associated with disease risk. *Genet Med.* 2019;0(0):1-9.
94. Stengel A, Kern W, Meggendorfer M, Haferlach T, Haferlach C. RUNX1 mutations in MDS, s-AML, and de novo AML: differences in accompanying genetic alterations and outcome. *Leuk Lymphoma.* 2019;60(5):1334-1336.
95. Drazer MW, Kadri S, Sukhanova M, et al. Prognostic tumor sequencing panels frequently identify germ line variants associated with hereditary hematopoietic malignancies. *Blood Adv.* 2018;2(2):146-150.
96. Kanagal-Shamanna R, Loghavi S, DiNardo CD, et al. Bone marrow pathologic abnormalities in familial platelet disorder with propensity for myeloid malignancy and germline RUNX1 mutation. *Haematologica.* 2017;102(10):1661-1670.
97. Bluteau D, Glembotsky AC, Raimbault A, et al. Dysmegakaryopoiesis of FPD/AML pedigrees with constitutional RUNX1 mutations is linked to myosin II deregulated expression. *Blood.* 2012;120(13):2708-2718.